



An Upper Bound on the Complexity of Recognizable Tree Languages

Olivier Finkel, Dominique Lecomte, Pierre Simonnet

► To cite this version:

Olivier Finkel, Dominique Lecomte, Pierre Simonnet. An Upper Bound on the Complexity of Recognizable Tree Languages. *RAIRO - Theoretical Informatics and Applications (RAIRO: ITA)*, 2015, 49 (2), pp.121-137. hal-01128609

HAL Id: hal-01128609

<https://hal.science/hal-01128609>

Submitted on 10 Mar 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

An Upper Bound on the Complexity of Recognizable Tree Languages

Olivier Finkel

Equipe de Logique Mathématique

Institut de Mathématiques de Jussieu - Paris Rive Gauche

CNRS et Université Paris Diderot Paris 7

Bâtiment Sophie Germain Case 7012

75205 Paris Cedex 13, France

E Mail: finkel@math.univ-paris-diderot.fr

Dominique Lecomte

Projet Analyse Fonctionnelle

Institut de Mathématiques de Jussieu - Paris Rive Gauche

Université Pierre et Marie Curie Paris 6

Couloir 16-26, 4ème étage, Case 247, 4, place Jussieu,

75252 Paris Cedex 05, France

E Mail: dominique.lecomte@upmc.fr

Université de Picardie, I.U.T. de l'Oise, site de Creil

13, allée de la faïencerie,

60107 Creil, France

Pierre Simonnet

UMR CNRS 6134

Faculté des Sciences

Université de Corse

Quartier Grossetti BP52 20250, Corte, France

E Mail: simonnet@univ-corse.fr

Abstract

The third author noticed in his 1992 PhD Thesis [Sim92] that every regular tree language of infinite trees is in a class $\mathfrak{D}(D_n(\Sigma_2^0))$ for some natural number $n \geq 1$, where \mathfrak{D} is the game quantifier. We first give a detailed exposition of this result. Next, using an embedding of the Wadge hierarchy of non self-dual Borel subsets of the Cantor space 2^ω into the class Δ_2^1 , and the notions of Wadge degree and Veblen function, we argue that this upper bound on the topological complexity of regular tree languages is much better than the usual Δ_2^1 .

keywords: infinite trees; tree automaton; regular tree language; Cantor topology; topological complexity; Borel hierarchy; game quantifier; Wadge classes; Wadge degrees; universal sets; provably- Δ_2^1 .

1 Introduction

A way to study the complexity of languages of infinite words or infinite trees accepted by various kinds of automata is to study their topological complexity, [LT94, PP04, GTW02, Sta97, Tho90, ADMN07]. In this paper we consider the topological complexity of regular languages of trees.

On one side the topological complexity of *deterministic* regular languages of trees has been determined. Niwinski and Walukiewicz proved that a tree language which is accepted by a deterministic Muller tree automaton is either in the class Π_3^0 or Π_1^1 -complete, [NW03]. And the Wadge hierarchy of *deterministic* regular languages of trees has been determined by Murlak, [Mur08, ADMN07].

On the other side in the course of years, more and more complex (non-deterministic) regular languages of trees have been found. Skurczyński proved that, for every natural number $n \geq 1$, there are some Π_n^0 -complete and some Σ_n^0 -complete regular tree languages, [Sku93]. Notice that it is an open question to know whether there exist some regular sets of trees which are Borel sets of infinite rank. Niwinski showed that there are some Σ_1^1 -complete regular sets of trees accepted by Büchi tree automata, and some Π_1^1 -complete regular sets of trees accepted by deterministic Muller tree automata, [Niw85]. Some examples of regular tree languages at some transfinite levels of the difference hierarchy of analytic sets were given by Simonnet and Finkel in [Sim92, FS09]. Hummel proved in [Hum12] that there exists some (unambiguous) regular tree language which is topologically more complex than any set in the difference hierarchy of analytic sets. Arnold and Niwinski showed in [AN08] that the game tree languages $W_{(\iota, \kappa)}$ form an infinite hierarchy of non Borel regular sets of trees with regard to the Wadge reducibility.

An upper bound on the complexity of regular languages of trees follows from the definition of acceptance by non deterministic Muller or Rabin automata and from Rabin's complementation Theorem: every regular set of trees is a Δ_2^1 -set, see [Rab69, PP04, Tho90, LT94].

The third author noticed in his 1992 PhD Thesis [Sim92] that every regular tree language of infinite trees is in a class $\mathfrak{D}(D_n(\Sigma_2^0))$ for some natural number $n \geq 1$, where \mathfrak{D} is the game quantifier. We first give a detailed exposition of this result. Next, using an embedding of the Wadge hierarchy of non self-dual Borel subsets of the Cantor space 2^ω into the class Δ_2^1 , and the notions of Wadge degree and Veblen function, we argue that this upper bound on the topological complexity of regular tree languages is much better than the usual Δ_2^1 .

The paper is organized as follows. In Section 2 we recall the notions of Muller tree automaton and regular tree language. The notions of topology are recalled in Section 3. We give the upper bound on the complexity of regular tree languages in Section 4. We argue that it is a much better upper bound than Δ_2^1 in Section 5.

2 Recognizable tree languages

We now recall usual notation of formal language theory.

- In the sequel, Σ will be a finite alphabet with at least two letters. A **non-empty finite word** over Σ is a sequence $x = a_1 \cdots a_k$, where $a_i \in \Sigma$ for $i = 1, \dots, k$, and $k \geq 1$ is a natural number. The **length** of x is k , and denoted by $|x|$. The **empty word** has no letter and is denoted by λ ; its length is 0. Σ^* is the **set of finite words** (including the empty word) over Σ . A **finitary language** over Σ is a subset of Σ^* .

- The **first infinite ordinal** is ω . An ω -**word** over Σ is an ω -sequence $a_1 \cdots a_n \cdots$, where $a_i \in \Sigma$ for each natural number $i \geq 1$. When σ is an ω -word over Σ , we write $\sigma = \sigma(1)\sigma(2) \cdots \sigma(n) \cdots$, where $\sigma(i) \in \Sigma$ for each i , $\sigma[n] = \sigma(1)\sigma(2) \cdots \sigma(n)$ for each $n \geq 1$, and $\sigma[0] = \lambda$. The **set of ω -words** over Σ is denoted by Σ^ω . An ω -**language** over Σ is a subset of Σ^ω .

- The usual concatenation product of two finite words u and v is denoted by $u \cdot v$ (and sometimes just uv). This product is extended to the product of a finite word u and an ω -word v : the infinite word $u \cdot v$ is then the ω -word such that: $(u \cdot v)(k) = u(k)$ if $k \leq |u|$, and $(u \cdot v)(k) = v(k - |u|)$ if $k > |u|$.

- The **prefix relation** is denoted by \sqsubseteq : a finite word u is a **prefix** of a finite word v (respectively, an infinite word v), denoted $u \sqsubseteq v$, if and only if there exists a finite word w (respectively, an infinite word w), such that $v = u \cdot w$.

- We now introduce the languages of infinite binary trees whose nodes are labelled in a finite alphabet Σ . A node of an infinite binary tree is represented by a finite word over the alphabet $\{l, r\}$, where l means “left” and r means “right”. An infinite binary tree whose nodes are labelled in Σ is identified with a function $t : \{l, r\}^* \rightarrow \Sigma$. The **set of infinite binary trees labelled in Σ** will be denoted by T_Σ^ω .

- Let t be a tree. A **branch** b of t is a subset of the set of nodes of t which is linearly ordered by the tree partial order \sqsubseteq and closed under prefix relation, i.e., if x and y are nodes of t such that $y \in b$ and $x \sqsubseteq y$, then $x \in b$. A branch b of a tree is said to be **maximal** if there is no branch of t containing strictly b .

Let t be an infinite binary tree in T_Σ^ω . If b is a maximal branch of t , then this branch is infinite. Let $(x_i)_{i \geq 0}$ be the enumeration of the nodes in b which is strictly increasing for the prefix order. The infinite sequence of labels of the nodes of such a maximal branch b , i.e., $t(x_0)t(x_1) \cdots t(x_n) \cdots$, is called a **path**. It is an ω -word over Σ .

We now define tree automata and regular tree languages.

Definition 2.1 A (nondeterministic topdown) **tree automaton** is a quadruple $\mathcal{A} = (Q, \Sigma, \Delta, q_0)$, where Q is a finite set of states, Σ is a finite input alphabet, $q_0 \in Q$ is the initial state and $\Delta \subseteq Q \times \Sigma \times Q \times Q$ is the transition relation. The tree automaton \mathcal{A} is said to be **deterministic** if the relation Δ is a functional one, i.e., if for each $(q, a) \in Q \times \Sigma$ there is at most one pair of states (q', q'') such that $(q, a, q', q'') \in \Delta$. A **run** of the tree automaton \mathcal{A} on an infinite binary tree $t \in T_\Sigma^\omega$ is a infinite binary tree $\rho \in T_Q^\omega$ such that $\rho(\lambda) = q_0$ and, for each $x \in \{l, r\}^*$, $(\rho(x), t(x), \rho(x \cdot l), \rho(x \cdot r)) \in \Delta$.

Definition 2.2 A **Muller** (nondeterministic topdown) **tree automaton** is a 5-tuple $\mathcal{A} = (Q, \Sigma, \Delta, q_0, \mathcal{F})$, where (Q, Σ, Δ, q_0) is a tree automaton and $\mathcal{F} \subseteq 2^Q$ is the collection of designated state sets. A run ρ of the Muller tree automaton \mathcal{A} on an infinite binary tree $t \in T_\Sigma^\omega$ is said to be **accepting** if, for each path p of ρ , the set of states appearing infinitely often in this path is in \mathcal{F} . The **tree language** $L(\mathcal{A})$ **accepted by the Muller tree automaton \mathcal{A}** is the set of infinite binary trees $t \in T_\Sigma^\omega$ such that there is (at least) one accepting run of \mathcal{A} on t . The class **REG** of **regular**, or **recognizable**, **tree languages** is the class of tree languages accepted by some Muller automaton.

Remark 2.3 A tree language is accepted by a Muller tree automaton iff it is accepted by some Rabin tree automaton iff it is accepted by some parity tree automaton. We refer for instance to [Tho90, GTW02, PP04, ADMN07] for the definition of Rabin tree automaton and of parity tree automaton.

3 Topology

We assume the reader to be familiar with basic notions of topology which may be found in [Mos09, LT94, Kec95, Sta97, PP04]. There is a natural metric on the set Σ^ω , which is called the **prefix metric** and defined as follows. For $\sigma, \sigma' \in \Sigma^\omega$ and $\sigma \neq \sigma'$ let $\delta(\sigma, \sigma') = 2^{-l_{\text{pref}(\sigma, \sigma')}$ where $l_{\text{pref}(\sigma, \sigma')}$ is the first natural number n such that the $(n+1)^{\text{st}}$ letter of σ is different from the $(n+1)^{\text{st}}$ letter of σ' . This metric induces on Σ^ω the usual topology for which **open** subsets of Σ^ω are of the form $W \cdot \Sigma^\omega$, where $W \subseteq \Sigma^*$. A set $L \subseteq \Sigma^\omega$ is **closed** if its complement $\Sigma^\omega \setminus L$ is an open set. The topological space Σ^ω is a Cantor space equipped with the usual Cantor topology.

- There is also a natural topology on the set T_Σ^ω [Mos09, LT94, Kec95]. It is defined by the following distance. Let t and s be two distinct infinite trees in T_Σ^ω . Then the distance between t and s is $\frac{1}{2^n}$ where n is the smallest natural number such that $t(x) \neq s(x)$ for some word $x \in \{l, r\}^*$ of length n . The open sets are then in the form $T_0 \cdot T_\Sigma^\omega$ where T_0 is a set of finite labelled trees. $T_0 \cdot T_\Sigma^\omega$ is the set of infinite binary trees which extend some finite labelled binary tree $t_0 \in T_0$, t_0 is here a sort of prefix, an “initial subtree” of a tree in $t_0 \cdot T_\Sigma^\omega$. It is well known that the set T_Σ^ω , equipped with this topology, is homeomorphic to the Cantor space 2^ω , hence also to the topological spaces Σ^ω , where Σ is finite.

- We now define the Borel Hierarchy of subsets of Σ^ω . It is defined similarly on the space T_Σ^ω .

Definition 3.1 *For a countable ordinal $\xi \geq 1$, the classes Σ_ξ^0 and Π_ξ^0 of the **Borel Hierarchy** on the topological space Σ^ω are defined as follows:*

Σ_1^0 *is the class of open subsets of Σ^ω ,*

Π_1^0 *is the class of closed subsets of Σ^ω ,*

and, for any countable ordinal $\xi \geq 2$,

Σ_ξ^0 *is the class of countable unions of subsets of Σ^ω in $\bigcup_{\eta < \xi} \Pi_\eta^0$,*

Π_ξ^0 *is the class of countable intersections of subsets of Σ^ω in $\bigcup_{\eta < \xi} \Sigma_\eta^0$.*

For a countable ordinal ξ , a subset of Σ^ω is a Borel set of **rank** ξ if it is in $\Sigma_\xi^0 \cup \Pi_\xi^0$ but not in $\bigcup_{\eta < \xi} (\Sigma_\eta^0 \cup \Pi_\eta^0)$.

- There exists another hierarchy beyond the Borel hierarchy, which is called the **projective hierarchy**. The classes Σ_n^1 and Π_n^1 of the projective hierarchy, defined for natural numbers $n \geq 1$, are obtained from the Borel hierarchy by successive applications of the operations of projection and complementation. The first level of the projective hierarchy is formed by the class Σ_1^1 of analytic sets and the class Π_1^1 of **co-analytic sets**, which are the complements of analytic sets.

In particular, the class of Borel subsets of Σ^ω is strictly contained in the class of analytic sets, which are obtained by projection of Borel sets.

Definition 3.2 *A subset A of Σ^ω is **analytic** if there is a Borel subset B of $(\Sigma \times 2)^\omega$, where $2 = \{0, 1\}$, such that $\sigma \in A \Leftrightarrow \exists \theta \in 2^\omega$ with $(\sigma, \theta) \in B$, where (σ, θ) is the infinite word over the alphabet $\Sigma \times 2$ such that $(\sigma, \theta)(i) = (\sigma(i), \theta(i))$ for each natural number $i \geq 1$.*

Remark 3.3 *In the above definition we could take B in the class Π_2^0 , [Mos09].*

The Borel hierarchy and the projective hierarchy on T_Σ^ω are defined from open sets in the same manner as in the case of the topological space Σ^ω .

- The notion of Wadge reducibility is defined via the reduction by continuous functions. In the sequel, Θ will be a finite alphabet with at least two letters. For $L \subseteq \Sigma^\omega$ and $L' \subseteq \Theta^\omega$, L is said to be **Wadge reducible** to L' , denoted by $L \leq_W L'$, if there exists a continuous function $f : \Sigma^\omega \rightarrow \Theta^\omega$, such that $L = f^{-1}(L')$. If $\xi \geq 1$ is a countable ordinal and $n \geq 1$ is a natural number, then a set $F \subseteq \Sigma^\omega$ is said to be Σ_ξ^0 (respectively, Π_ξ^0 , Σ_n^1 , Π_n^1)-**complete** if, for any set $E \subseteq \Theta^\omega$, $E \in \Sigma_\xi^0$ (respectively, $E \in \Pi_\xi^0$, $E \in \Sigma_n^1$, $E \in \Pi_n^1$) iff $E \leq_W F$.

The ω -language $\mathcal{R} = (0^* \cdot 1)^\omega$ is a well known example of a Π_2^0 -complete subset of $\{0, 1\}^\omega$. It is the set of ω -words over $\{0, 1\}$ having infinitely many occurrences of the letter 1. Its complement $\{0, 1\}^\omega \setminus (0^* \cdot 1)^\omega$ is a Σ_2^0 -complete subset of $\{0, 1\}^\omega$. The set of infinite trees in T_Σ^ω , where $\Sigma = \{0, 1\}$, having at least one path in the ω -language \mathcal{R} is Σ_1^1 -complete.

- We now define the difference hierarchy over a class Γ . Let n be a natural number, and $(A_p)_{p < n}$ be an increasing sequence of subsets of some space Z . The set $D_n[(A_p)_{p < n}]$ is the set of elements $z \in Z$ such that $z \in A_p \setminus \bigcup_{q < p} A_q$ for some $p < n$ whose parity is opposite to that of n . We can now define the class of n -differences of Γ -subsets of Z , where $Z = \Sigma^\omega$ or $Z = T_\Sigma^\omega$:

$$D_n(\Gamma) := \{D_n[(A_p)_{p < n}] \mid A_p \text{ is in the class } \Gamma \text{ for each } p < n\}.$$

It is well known that, for every countable ordinal $\xi \geq 1$, the hierarchy of differences of Σ_ξ^0 -sets is strict, i.e., the inclusion $D_m(\Sigma_\xi^0) \subset D_n(\Sigma_\xi^0)$ holds if $m < n$.

Notice that the difference hierarchy has an extension to countable ordinal ranks, see for instance [Mos09, Kec95], but we shall not need this in the sequel.

Moreover, every regular ω -language is a boolean combination of Σ_2^0 -sets, and belongs to some class $D_n(\Sigma_2^0)$, for some natural number $n \geq 1$. We shall also consider these classes $D_n(\Sigma_2^0)$ in the sequel.

It is also well known that the hierarchy of differences of analytic sets is strict. It was proved by Hummel in [Hum12] that there exists some (unambiguous) regular tree language which is topologically more complex than any set in the difference hierarchy of analytic sets because there exists some (unambiguous) regular tree language which does not belong to the σ -algebra generated by the analytic sets.

4 An Upper Bound

We first recall the definition of Gale-Stewart games.

Definition 4.1 ([Kec95]) *Let $A \subseteq \Sigma^\omega$. The **Gale-Stewart game** $G(A)$ is a game with perfect information between two players. Player 1 first writes a letter $a_1 \in \Sigma$, then Player 2 writes a letter $b_1 \in \Sigma$, then Player 1 writes $a_2 \in \Sigma$, and so on \dots . After ω steps, the two players have composed a word $\sigma = a_1b_1a_2b_2\dots$ of Σ^ω . Player 1 wins the play iff $\sigma \in A$, otherwise Player 2 wins the play.*

*Let $A \subseteq \Sigma^\omega$ and $G(A)$ be the associated Gale-Stewart game. A **strategy** for Player 1 is a function $F_1 : (\Sigma^2)^* \rightarrow \Sigma$ and a strategy for Player 2 is a function $F_2 : (\Sigma^2)^* \Sigma \rightarrow \Sigma$. Player 1 follows the strategy F_1 in a play if for each natural number $n \geq 1$ $a_n = F_1(a_1b_1a_2b_2\dots a_{n-1}b_{n-1})$. If Player 1 wins every play in which she has followed the strategy F_1 , then we say that the strategy F_1 is a **winning strategy** (w.s.) for Player 1. The notion of winning strategy for Player 2 is defined in a similar manner.*

*The game $G(A)$ is said to be **determined** if one of the two players has a winning strategy.*

We now recall the definition of the **game quantifier**.

Definition 4.2 ([Mos09]) (1) *If $P \subseteq \Sigma^\omega \times 2^\omega$, then we define*

$$\mathcal{D}P := \left\{ \sigma \in \Sigma^\omega \mid \text{Player 1 has a winning strategy in the game } G(P_\sigma) \right\}$$

where $P_\sigma = \{ \alpha \in 2^\omega \mid (\sigma, \alpha) \in P \}$.

(2) *If $\mathbf{\Gamma}$ is a class of sets and $L \subseteq \Sigma^\omega$, then $L \in \mathcal{D}\mathbf{\Gamma}$ if and only if there is $P \subseteq \Sigma^\omega \times 2^\omega$ in $\mathbf{\Gamma}$ such that $L = \mathcal{D}P$.*

Note that one can replace Σ^ω with the homeomorphic space T_Σ^ω in this definition, replacing also P with a $\mathbf{\Gamma}$ -subset of $T_\Sigma^\omega \times 2^\omega$. We obtain like this the definition of subsets of T_Σ^ω in the class $\mathcal{D}(\mathbf{\Gamma})$. Note that we consider the alphabet $2 = \{0, 1\}$ and not the finite alphabets with at least two letters Σ in order to avoid troubles in the proof of Proposition 5.8 to come. This leads us to prove the following lemma.

Lemma 4.3 *We can find a continuous injection $\varphi: \Sigma^\omega \rightarrow 2^\omega$ and an open subset O of 2^ω such that, for each subset A of Σ^ω , the set $A' := \varphi[A] \cup O$ satisfies the following property: Player 1 has a winning strategy in the game $G(A)$ if and only if Player 1 has a winning strategy in the game $G(A')$.*

Proof. The proof is similar to that of Lemma 3.14 in [Fin13b]. We may assume that Σ is a natural number and with a usual convention that $\Sigma = \{0, 1, \dots, \Sigma - 1\}$. Let $\varphi: \Sigma^\omega \rightarrow 2^\omega$ be defined by $\varphi(a_1 a_2 \dots) := (11)^{a_1} 0 (11)^{a_2} 0 \dots$. Note that φ is a continuous injection. We set $D_1 := \varphi[A]$, so that D_1 codes A . We also set

$$\begin{aligned} D_2 &:= \{\alpha \in 2^\omega \mid \exists a_1, \dots, a_{2n} \in \Sigma \ \exists k < \Sigma - 1 \ (11)^{a_1} 0 \dots (11)^{a_{2n}} 0 1^{2k+1} 0 \sqsubseteq \alpha\}, \\ D_3 &:= \{\alpha \in 2^\omega \mid \exists a_1, \dots, a_{2n+1} \in \Sigma \ (11)^{a_1} 0 \dots (11)^{a_{2n+1}} 0 1^{2\Sigma-1} \sqsubseteq \alpha\}. \end{aligned}$$

As D_2 and D_3 are open, $O := D_2 \cup D_3$ is open too. The sets D_2, D_3 are the results of infinite plays where two players alternatively write a letter 0 or 1 and the infinite word written by the two players in ω steps is out of $K := \varphi[\Sigma^\omega]$, due to the letters written by Player 2. More precisely, D_2 is the set of plays where Player 2 writes the $(2n+1)$ th letter 0 while it was Player 1's turn to do this. And D_3 is the set of plays where Player 2 does not write a letter 0 soon enough when it is his turn to do this.

If now the two players write alternatively a letter 0 or 1 in such a way that the infinite word written by them in ω steps is in K and of the form $(11)^{a_1} 0 (11)^{a_2} 0 \dots$, then the letters 0 have been written alternatively by Player 1 and by Player 2, and these letters 0 determine the natural numbers a_i . Thus the natural numbers a_{2i+1} have been chosen by Player 1, and the natural numbers a_{2i+2} have been chosen by Player 2 (for $i \in \omega$).

If Player 1 has a winning strategy s in the game $G(A)$, then consider a play in the game $G(A')$. If the two players alternatively write a letter 0 or 1 and the infinite word α written by them in ω steps is in K and of the form $(11)^{a_1} 0 (11)^{a_2} 0 \dots$, then the game is like a game where each player writes a letter in Σ at each step of the play, and Player 1 can apply s to ensure that $(a_i)_{i \geq 1} \in A$, which implies that $\varphi((a_i)_{i \geq 1}) \in D_1 \subseteq A'$, so Player 1 wins the play. If α is out of K due to the letters written by Player 2, then $\alpha \in D_2 \cup D_3 \subseteq A'$, and Player 1 wins the play again. This shows that Player 1 has a winning strategy in $G(A')$.

If Player 1 has a winning strategy s' in the game $G(A')$, then consider a play in the game $G(A)$ where Player 2 does not make the final word α in $D_2 \cup D_3$. Player 1, following s' , must write letters so that the final word α belongs to K . Then the game is reduced to the game $G(A)$ in which the two players alternatively write letters a_i in Σ . But Player 1 wins the game and this implies that Player 1 has a winning strategy in $G(A)$. \square

We now give the upper bound on the complexity of regular tree languages.

Theorem 4.4 *Let \mathcal{A} be a Muller tree automaton. Then the tree language $L(\mathcal{A})$ is in the class $\mathfrak{D}(D_n(\Sigma_2^0))$, for some natural number $n \geq 1$.*

Proof. Let $\mathcal{A} = (Q, \Sigma, \Delta, q_0, \mathcal{F})$ be a Muller tree automaton, where (Q, Σ, Δ, q_0) is a tree automaton and $\mathcal{F} \subseteq 2^Q$ is the collection of designated state sets.

The idea is to use a game which was considered by Gurevich and Harrington in [GH82]. For $t \in T_\Sigma^\omega$ we consider the Gale-Stewart game $G(T_t)$, where the set $T_t \subseteq (\Delta \cup \{l, r\})^\omega$ is defined as follows. Intuitively, in the game $G(T_t)$, Player 1 writes transitions of the automaton \mathcal{A} , i.e., letters of Δ , and Player 2 writes letters l or r , so that at the end of a play he has actually composed a path in the tree t .

At the first step, Player 1 chooses a transition $(q_0, t(\lambda), q, q') \in \Delta$. Next, Player 2 chooses l or r . If Player 2 chooses l this means he has chosen the left son of the root node of the tree and thus Player 1 has to choose a transition of the form $(q, t(l), q'', q''')$. If Player 2 chooses r this means he has chosen the right son of the root node of the tree and thus Player 1 has to choose a transition of the form $(q', t(r), q'', q''')$. The game continues like this and Player 1 wins this game if the set of states (played by Player 1) which appear infinitely often on the path chosen by Player 2 belongs to \mathcal{F} .

Formally, if $\alpha \in \Theta^\omega$, then we define $(\alpha)_1, (\alpha)_2 \in \Theta^\omega$ by $(\alpha)_1(k) := \alpha(2k - 1)$ and $(\alpha)_2(k) := \alpha(2k)$. T_t is the union of $\left\{ \alpha \in (\Delta \cup \{l, r\})^\omega \mid \exists k \geq 1 (\alpha)_2(k) \notin \{l, r\} \right\}$ with

$$\left\{ \alpha \in (\Delta \cup \{l, r\})^\omega \mid \forall k \geq 1 \left((\alpha)_1(k) \in \Delta \text{ and } (\alpha)_2(k) \in \{l, r\} \text{ and } \alpha(1)(1) = q_0 \right. \right. \\ \left. \text{and } (\alpha)_1(k)(2) = t((\alpha)_2[k - 1]) \text{ and } (\alpha)_1(k + 1)(1) = \begin{cases} (\alpha)_1(k)(3) & \text{if } (\alpha)_2(k) = l \\ (\alpha)_1(k)(4) & \text{if } (\alpha)_2(k) = r \end{cases} \right. \\ \left. \text{and } \{q \in Q \mid \forall j \in \omega \exists m \geq j (\alpha)_1(m)(1) = q\} \in \mathcal{F} \right\}.$$

By definition of the Muller acceptance condition for the tree automaton \mathcal{A} ,

$$t \in L(\mathcal{A}) \Leftrightarrow \text{Player 1 has a winning strategy in the game } G(T_t).$$

Indeed, if $t \in L(\mathcal{A})$, then there is a witness $\rho \in T_Q^\omega$ and we can define a strategy $\tilde{\rho}$ for Player 1 by $\tilde{\rho}(x) := (\rho(x), t(x), \rho(xl), \rho(xr))$, and $\tilde{\rho}$ is winning for Player 1. Conversely, if $\tilde{\rho}$ is a winning strategy for Player 1, then the formula $\rho(x) := \tilde{\rho}(x)(1)$ defines a witness for the fact that $t \in L(\mathcal{A})$.

We apply Lemma 4.3 to the finite alphabet $\Delta \cup \{l, r\}$, which gives a continuous injection $\varphi: (\Delta \cup \{l, r\})^\omega \rightarrow 2^\omega$ and an open subset O of 2^ω . Then we set

$$P := \{(t, \alpha) \in T_\Sigma^\omega \times 2^\omega \mid \alpha \in \varphi[T_t] \cup O\},$$

so that $t \in L(\mathcal{A}) \Leftrightarrow \text{Player 1 has a winning strategy in the game } G(P_t) \Leftrightarrow t \in \mathfrak{D}P$.

It remains to see that P is in $D_n(\Sigma_2^0)$ for some natural number $n \geq 1$. If $B := D_n[(A_p)_{p < n}]$ is in $D_n(\Sigma_2^0)$ and C is a closed set, then $B \cap C = D_n[(A_p \cap C)_{p < n}]$ is also in $D_n(\Sigma_2^0)$. If U is an open set, then $B \cup U = D_n[(A_p \cup U)_{p < n}]$ if n is odd, and $B \cup U = D_n[(A_0 \setminus U, A_1 \cup U, \dots, A_{n-1} \cup U)]$ if n is even, so that $B \cup U$ is also in $D_n(\Sigma_2^0)$. This shows that the class $D_n(\Sigma_2^0)$ is closed under intersections with a closed set and unions with an open set. As it is also closed under images by homeomorphisms and φ is a homeomorphism onto its compact range, it is enough to show that T is in $D_n(\Sigma_2^0)$ for some natural number $n \geq 1$. This is known to be equivalent to the statement “ T is a boolean combination of Σ_2^0 -sets”, and this follows from the definition of the Muller acceptance condition of the tree automaton \mathcal{A} . Indeed,

- the condition “ $\exists k \geq 1 (\alpha)_2(k) \notin \{l, r\}$ ” is open,
- the condition involving k and the big parentheses in the definition of T_t is closed,
- the condition involving \mathcal{F} says that there is F in the finite set \mathcal{F} such that, for each q in the finite set Q , either $q \in F$ and $\forall j \in \omega \exists m \geq j (\alpha)_1(m)(1) = q$ (this is a Π_2^0 condition), or $q \notin F$ and $\exists j \in \omega \forall m \geq j (\alpha)_1(m)(1) \neq q$ (this is a Σ_2^0 condition). This finishes the proof. \square

5 The Upper Bound is much better than Δ_2^1

In this section, we are going to show that the upper bound given in the preceding section is actually much better than the usual one, Δ_2^1 .

- We first define the Wadge hierarchy, which is a great refinement of the Borel hierarchy, and is defined via reductions by continuous functions, [Dup01, Wad83].

Definition 5.1 (Wadge [Wad83]) *Let $L \subseteq \Sigma^\omega$ and $L' \subseteq \Theta^\omega$. We say that L and L' are **Wadge equivalent** if $L \leq_W L'$ and $L' \leq_W L$. This will be denoted by $L \equiv_W L'$. We shall also say that $L <_W L'$ iff $L \leq_W L'$ and $L' \not\leq_W L$. We say that L is **self dual** if $L \equiv_W \neg L$ ($\neg L$ is the complement of L), otherwise we say that L is **non self dual**.*

The relation \leq_W is reflexive and transitive, and \equiv_W is an equivalence relation. The equivalence classes of \equiv_W are called **Wadge degrees**. The **Wadge hierarchy** WH is the class of Borel subsets of a space Σ^ω , equipped with \leq_W and \equiv_W . For $L \subseteq \Sigma^\omega$ and $L' \subseteq \Theta^\omega$, if $L \leq_W L'$ and $L = f^{-1}(L')$, where f is a continuous function from Σ^ω into Θ^ω , then f is called a **continuous reduction** of L to L' . Intuitively, it means that L is less complicated than L' , because in order to check whether $\sigma \in L$, it suffices to check whether $f(\sigma) \in L'$ (where f is a continuous function). Hence the Wadge degree of an ω -language is a measure of its topological complexity.

Note that in the above definition, we consider that a subset $L \subseteq \Sigma^\omega$ is given together with the alphabet Σ . This is important as it is shown by the following simple example. Let $L_1 = \{0, 1\}^\omega \subseteq \{0, 1\}^\omega$ and $L_2 = \{0, 1\}^\omega \subseteq \{0, 1, 2\}^\omega$. So the languages L_1 and L_2 are equal but considered over the different alphabets $\Sigma_1 = \{0, 1\}$ and $\Sigma_2 = \{0, 1, 2\}$. It turns out that $L_1 <_W L_2$. In fact L_1 is open *and* closed in Σ_1^ω while L_2 is closed but not open in Σ_2^ω .

We can now define the Wadge class of a set L .

Definition 5.2 *Let L be a subset of Σ^ω . The **Wadge class** of L is*

$$[L] = \{L' \mid L' \subseteq \Theta^\omega \text{ for a finite alphabet } \Theta \text{ and } L' \leq_W L\}.$$

Recall that each Borel class Σ_ξ^0 and Π_ξ^0 is a Wadge class. It follows from the study of the Wadge hierarchy that a set $L \subseteq X^\omega$ is Σ_ξ^0 (respectively, Π_ξ^0)-complete iff it is in Σ_ξ^0 but not in Π_ξ^0 (respectively, in Π_ξ^0 but not in Σ_ξ^0).

Using the notion of a Wadge game, and the fact that the determinacy of Wadge games follows from Martin's Theorem stating that every Gale-Stewart Game $G(B)$, with B a Borel set, is determined, see [Kec95], Wadge proved the following result.

Theorem 5.3 (Wadge) *Up to complement and \equiv_W , the class of Borel subsets of the spaces Σ^ω is a well ordered hierarchy. We can find an ordinal $|WH|$, called the length of the hierarchy, and a map d_W^0 from WH onto $|WH| \setminus \{0\}$, such that, for all $L, L' \subseteq \Sigma^\omega$,*

- $d_W^0 L < d_W^0 L' \Leftrightarrow L <_W L'$,
- $d_W^0 L = d_W^0 L' \Leftrightarrow [L \equiv_W L' \text{ or } L \equiv_W \neg L']$.

• Let ω_1 be the **first uncountable ordinal**. The Wadge hierarchy of Borel sets of finite rank has length ${}^1\varepsilon_0$ where ${}^1\varepsilon_0$ is the limit of the ordinals α_n defined by $\alpha_1 = \omega_1$ and $\alpha_{n+1} = \omega_1^{\alpha_n}$ if n is a natural number. Then ${}^1\varepsilon_0$ is the first fixed point of the ordinal exponentiation of base ω_1 . The length of the Wadge hierarchy of Borel sets in $\Delta_\omega^0 = \Sigma_\omega^0 \cap \Pi_\omega^0$ is the ω_1^{th} fixed point of the ordinal exponentiation of base ω_1 , which is a much larger ordinal. The length of the whole Wadge hierarchy is described in [Wad83, Dup01], and uses the Veblen functions. To recall the definition of these functions, we need the notion of cofinality of an ordinal which may be found in [Jec02] and which we briefly recall now.

Definition 5.4 *Let ζ be a limit ordinal. The **cofinality** of ζ , denoted by $\text{cof}(\zeta)$, is the least ordinal β such that there exists a strictly increasing sequence of ordinals $(\zeta_i)_{i < \beta}$, of length β , such that $\sup_{i < \beta} \zeta_i = \zeta$ and $\zeta_i < \zeta$ for each $i < \beta$. This definition is usually extended to 0 and to the successor ordinals: $\text{cof}(0) = 0$ and $\text{cof}(\zeta + 1) = 1$ for every ordinal ζ .*

The cofinality of a limit ordinal is always a limit ordinal with $\omega \leq \text{cof}(\zeta) \leq \zeta$. The ordinal $\text{cof}(\zeta)$ is in fact a cardinal (see [Jec02]). If the cofinality of a limit ordinal ζ is $\leq \omega_1$, then only the following cases may happen: $\text{cof}(\zeta) = \omega$ or $\text{cof}(\zeta) = \omega_1$. In the sequel we do not need to consider cofinalities which are larger than ω_1 . In the sequel, ω_2 will be the first ordinal of cardinality greater than \aleph_1 , the cardinal of ω_1 . Note that each ordinal $\zeta < \omega_2$ has cofinality smaller than or equal to ω_1 .

Definition 5.5 *The Veblen hierarchy* $(V_\xi)_{\xi < \omega_1}$ of functions from $\omega_2 \setminus \{0\}$ into itself is defined as follows. V_0 is the function that enumerates ordinals of cofinality ω_1 or 1 that are closed under ordinal addition, i.e., $V_0(1) = 1$, $V_0(\zeta + 1) = V_0(\zeta) \cdot \omega_1$, $V_0(\zeta) = \omega_1^\zeta$ when $\text{cof}(\zeta) = \omega_1$, $V_0(\zeta) = \omega_1^{\zeta+1}$ when $\text{cof}(\zeta) = \omega$. For $\xi > 0$, V_ξ is the function that enumerates ordinals of cofinality ω_1 or 1 that are closed under each function V_η for any $\eta < \xi$.

We can now describe the length of the whole Wadge hierarchy of Borel sets on a Cantor space Σ^ω or T_Σ^ω . It is the ordinal $\sup_{\xi < \omega_1} V_\xi(2)$. This is really a huge ordinal, with regard to the ω_1^{th} fixed point of the ordinal exponentiation of base ω_1 , which is the length of the Wadge hierarchy of Borel sets in Δ_ω^0 .

- We now recall the notion of a universal set which will be useful in the sequel.

Definition 5.6 *Let Γ be a class of sets. We say that $\mathcal{U} \subseteq 2^\omega \times \Sigma^\omega$ is **universal** for the Γ subsets of Σ^ω if \mathcal{U} is in Γ , and for each $A \subseteq \Sigma^\omega$ in Γ there is $\beta \in 2^\omega$ such that A is the vertical section $\mathcal{U}_\beta := \{\alpha \in \Sigma^\omega \mid (\beta, \alpha) \in \mathcal{U}\}$ of \mathcal{U} at β .*

The following result is mentioned in [LSR88].

Theorem 5.7 *Let Γ be the Wadge class of a non self-dual Borel subset of 2^ω . Then there is a universal set for the Γ subsets of Σ^ω .*

Proof. Let $A, B \subseteq 2^\omega$. We consider the **Wadge game** $G(A, B)$ associated with A and B . It is the Gale-Stewart game defined by

$$\text{Player 2 wins the play} \Leftrightarrow (a_1 a_2 \dots \in A \Leftrightarrow b_1 b_2 \dots \in B).$$

If s is a strategy for Player 2 and $\alpha \in 2^\omega$, then we denote by $s * \alpha$ the element $b_1 b_2 \dots$ of 2^ω given by the answers of Player 2 in the play where Player 1 plays $\alpha := a_1 a_2 \dots$ and Player 2 follows s . Formally, $(s * \alpha)(1) := b_1 := s(a_1)$ and, inductively, $(s * \alpha)(k) := b_k := s(a_1 b_1 a_2 \dots b_{k-1} a_k)$ if $k \geq 2$. Notice that a strategy s for Player 2 is an element of $2^{(2^2)^*2}$. As $(2^2)^*2$ is countable, we can identify $2^{(2^2)^*2}$ with the Cantor space 2^ω . Note that the map $e: (s, \alpha) \mapsto s * \alpha$ is continuous from $2^{(2^2)^*2} \times 2^\omega$ into 2^ω .

Theorem 2.7 and Remark 4 after the proof of Theorem 3.1 in [LSR88] provide a subset L of 2^ω in $\mathbf{\Gamma}$ such that Player 2 has a winning strategy in the game $G(A, L)$ for each subset A of 2^ω in $\mathbf{\Gamma}$. We set $\mathcal{V} := \{(s, \alpha) \in 2^{(2^2)^*2} \times 2^\omega \mid s * \alpha \in L\}$. As e is continuous, \mathcal{V} is in $\mathbf{\Gamma}$. If $A \subseteq 2^\omega$ is in $\mathbf{\Gamma}$, then Player 2 wins the Wadge game $G(A, L)$, which gives s in $2^{(2^2)^*2}$ such that $A = \mathcal{V}_s$. This shows that we can consider \mathcal{V} as universal for the $\mathbf{\Gamma}$ subsets of 2^ω , up to identification.

Let Σ be a finite alphabet, and $\psi : \Sigma^\omega \rightarrow 2^\omega$ be a homeomorphism. We set $\mathcal{U} := (\text{Id}_{2^\omega} \times \psi)^{-1}(\mathcal{V})$, and it is routine to check that \mathcal{U} is universal for the $\mathbf{\Gamma}$ subsets of Σ^ω . \square

Proposition 5.8 *Let $\mathbf{\Gamma}$ be a class of sets. If there is a universal for the $\mathbf{\Gamma}$ subsets of $\Sigma^\omega \times 2^\omega$, then there is a universal for the $\mathbf{\Gamma}$ subsets of Σ^ω .*

Proof. Let \mathcal{U} be a universal for the $\mathbf{\Gamma}$ subsets of $\Sigma^\omega \times 2^\omega$. We set $\mathcal{V} := \mathcal{U}$. Then $\mathcal{V} \subseteq 2^\omega \times \Sigma^\omega$ is in $\mathbf{\Gamma}$. Let $A \subseteq \Sigma^\omega$ be in $\mathbf{\Gamma}$, and $P \subseteq \Sigma^\omega \times 2^\omega$ in $\mathbf{\Gamma}$ such that $A = \mathcal{V}_P$. Then there is $\beta \in 2^\omega$ with $P = \mathcal{U}_\beta$. It remains to note that $A = \mathcal{V}_\beta$. \square

Corollary 5.9 *Let $\mathbf{\Gamma}$ be the Wadge class of a non self-dual Borel subset of 2^ω . Then there is a universal for the $\mathbf{\Gamma}$ subsets of Σ^ω .*

Proof. Theorem 5.7 gives a universal for the $\mathbf{\Gamma}$ subsets of $\Sigma^\omega \times 2^\omega$, and Proposition 5.8 gives the result. \square

• We now turn to the key result of embeddability of the Wadge hierarchy into that obtained with the game quantifier.

Lemma 5.10 *Let $\mathbf{\Gamma}$ be a class of sets closed under continuous pre-images. Then $\mathbf{\Gamma}$ is also closed under continuous pre-images.*

Proof. Let $L \subseteq \Sigma^\omega$ in $\mathbf{\Gamma}$, $P \subseteq \Sigma^\omega \times 2^\omega$ in $\mathbf{\Gamma}$ such that $L = \mathcal{V}_P$, and $f : \Theta^\omega \rightarrow \Sigma^\omega$ be continuous. As $\mathbf{\Gamma}$ is closed under continuous pre-images, the set

$$Q := \{(\delta, \alpha) \in \Theta^\omega \times 2^\omega \mid (f(\delta), \alpha) \in P\}$$

is in $\mathbf{\Gamma}$. Note that, for each $\delta \in \Theta^\omega$,

$$\begin{aligned} \delta \in f^{-1}(L) &\Leftrightarrow f(\delta) \in \mathcal{V}_P \\ &\Leftrightarrow \text{Player 1 has a winning strategy in the game} \\ &\quad G(\{\alpha \in 2^\omega \mid (f(\delta), \alpha) \in P\}) \\ &\Leftrightarrow \text{Player 1 has a winning strategy in the game } G(\{\alpha \in 2^\omega \mid (\delta, \alpha) \in Q\}) \\ &\Leftrightarrow \delta \in \mathcal{V}_Q. \end{aligned}$$

Thus $f^{-1}(L) \subseteq \Theta^\omega$ is in $\mathbf{\Gamma}$. \square

Corollary 5.11 *The map $\Gamma \mapsto \mathfrak{D}\Gamma$, defined on the collection of Wadge classes of a non self-dual Borel subset of 2^ω , is strictly increasing for the inclusion.*

Proof. Let L, L' be non self-dual Borel subsets of 2^ω , and Γ, Γ' be the Wadge classes of L, L' respectively. We assume that $\Gamma \subsetneq \Gamma'$, so that $L <_W L'$. By Wadge's lemma, $\neg L \leq_W L'$, so that Γ and $\check{\Gamma} := \{\neg A \mid A \in \Gamma\}$ are contained in Γ' (see 21.14 in [Kec95]). This implies that $\mathfrak{D}\Gamma \subseteq \mathfrak{D}\Gamma'$, and in fact $\mathfrak{D}\Gamma \cup \mathfrak{D}\check{\Gamma} \subseteq \mathfrak{D}\Gamma'$.

Assume, towards a contradiction, that $\mathfrak{D}\Gamma = \mathfrak{D}\Gamma'$. Corollary 5.9 gives a universal \mathcal{U} for the $\mathfrak{D}\Gamma'$ subsets of 2^ω . We set $A := \{\alpha \in 2^\omega \mid (\alpha, \alpha) \notin \mathcal{U}\}$. As \mathcal{U} is in $\mathfrak{D}\Gamma'$, it is also in $\mathfrak{D}\Gamma$. This gives $P \subseteq (2^\omega)^3$ in Γ with $\mathcal{U} = \mathfrak{D}P$. As L is Borel, 20.5 in [Kec95] implies that

$$\begin{aligned} (\beta, \gamma) \notin \mathcal{U} &\Leftrightarrow \neg(\text{Player 1 has a winning strategy in } G(\{\alpha \in 2^\omega \mid (\beta, \gamma, \alpha) \in P\})) \\ &\Leftrightarrow \text{Player 2 has a winning strategy in } G(\{\alpha \in 2^\omega \mid (\beta, \gamma, \alpha) \notin P\}). \end{aligned}$$

By 6D.1 in [Mos09], $\neg \mathcal{U} \in \mathfrak{D}\check{\Gamma} \subseteq \mathfrak{D}\Gamma'$. As the map $\alpha \mapsto (\alpha, \alpha)$ is continuous, $A \in \mathfrak{D}\Gamma'$, by Lemma 5.10. This gives $\beta \in \omega^\omega$ such that $A = \mathcal{U}_\beta$. Thus

$$(\beta, \beta) \in \mathcal{U} \Leftrightarrow \beta \in A \Leftrightarrow (\beta, \beta) \notin \mathcal{U},$$

which is absurd. □

Lemma 5.12 *Let Γ be a class of Borel sets. Then $\mathfrak{D}\Gamma$ is contained in Δ_2^1 .*

Proof. Let $L \subseteq \Sigma^\omega$ be in $\mathfrak{D}\Gamma$, and $P \subseteq \Sigma^\omega \times 2^\omega$ be in Γ such that $L = \mathfrak{D}P$. If σ, τ are strategies for Players 1 and 2 respectively, then we denote by $\sigma * \tau$ the element of 2^ω obtained when Player 1 follows σ and Player 2 follows τ . As P is Borel, $P_\beta := \{\alpha \in 2^\omega \mid (\beta, \alpha) \in P\}$ is Borel for each $\beta \in \Sigma^\omega$. By 20.5 in [Kec95], the game defined by P_β is determined. This implies that $\beta \in L$ iff

$$\begin{aligned} \beta \in \mathfrak{D}P &\Leftrightarrow \text{Player 1 has a winning strategy in the game } G(\{\alpha \in 2^\omega \mid (\beta, \alpha) \in P\}) \\ &\Leftrightarrow \exists \sigma \forall \tau (\beta, \sigma * \tau) \in P \Leftrightarrow \forall \tau \exists \sigma (\beta, \sigma * \tau) \in P. \end{aligned}$$

Thus $L \in \Delta_2^1$. □

The classes $D_n(\Sigma_2^0)$ appearing in Theorem 4.4 are contained in Σ_3^0 , which is a small part of the class of Borel sets. By Theorem 4.4 and Corollary 5.11, the tree languages of Muller tree automata are in the class $\mathfrak{D}\Sigma_3^0$. By 22.4 in [Kec95], the sequence $(\Sigma_\xi^0)_{\xi < \omega_1}$ is strictly increasing for the inclusion. By Corollary 5.11, the sequence $(\mathfrak{D}\Sigma_\xi^0)_{\xi < \omega_1}$ is also strictly increasing for the inclusion. By Lemma 5.12, the elements of the latter sequence are contained in the class Δ_2^1 . This shows already that the increasing sequence $(\mathfrak{D}(D_n(\Sigma_2^0)))_{n \geq 1}$ is only a small part of the hierarchy of Δ_2^1 sets.

This is actually emphasized by considering the Wadge hierarchy of non self-dual Borel subsets of 2^ω . By Corollary 5.11 we know that this hierarchy can be embedded into a hierarchy of classes $\mathfrak{D}\Gamma$, included into Δ_2^1 . On the other hand the Borel class Σ_2^0 is known to be a non self-dual Wadge class and the Wadge degree (in the sense of Theorem 5.3) of any Σ_2^0 -complete set is equal to ω_1 . Moreover, for each natural number $n \geq 1$, the class $D_n(\Sigma_2^0)$ is also a non self-dual Wadge class and the Wadge degree of any $D_n(\Sigma_2^0)$ -complete is equal to $(\omega_1)^n$, [Wad83, Dup01]. Therefore, we see that this ordinal is actually much smaller than the first fixed point of the ordinal exponentiation of base ω_1 (the length of the Wadge hierarchy of Borel sets of finite ranks), than the ω_1^{th} fixed point of the ordinal exponentiation of base ω_1 (the length of the Wadge hierarchy of Borel sets in Δ_ω^0), and a fortiori than the ordinal

$$\sup_{\xi < \omega_1} V_\xi(2),$$

which is the length of the Wadge hierarchy of (non self-dual) Borel subsets of the Baire space or the Cantor space. In conclusion the upper bound we gave is much smaller than Δ_2^1 , as summarized in the following theorem.

Theorem 5.13 *The hierarchy of classes $\mathfrak{D}\Gamma$, for non self-dual Borel Wadge classes Γ of 2^ω , is a hierarchy of length $\sup_{\xi < \omega_1} V_\xi(2)$ included in the class Δ_2^1 , while the regular tree languages are contained in the first $(\omega_1)^\omega$ levels of this hierarchy.*

- We also note that regular tree languages are in the class provably- Δ_2^1 (see [Kan97, page 180]). This is essentially proved in [HKMN08] from Rabin's Theorem [Rab69]. Thus all regular tree languages have the Baire property (see [Kan97, page 180]). This gives an answer to a question of the third author in [Sim92].

As noticed by the referee of this paper, this also follows from [GMMS14], since all R-sets are Baire-measurable as well as Lebesgue measurable and it is proved in [GMMS14] that regular tree languages are R-sets.

6 Concluding remarks

We gave an upper bound on the topological complexity of recognizable tree languages and showed, using an embedding of the Wadge hierarchy of non self-dual Borel sets of 2^ω , that this upper bound is actually much smaller than Δ_2^1 .

The anonymous referee of this paper indicated us that since the first appearance in 1992 of the above Theorem 4.4 in the PhD Thesis of the third author of this paper, quite a few authors made various observations in a few papers that we now mention.

In J. Bradfield’s paper [Bra03], and in [BDQ05] by J. Bradfield, J. Duparc and S. Quickert, a link between the game quantifier and the μ -calculus is described, and Corollary 11 of [Bra03] explicitly states an upper bound in a style similar to the one presented in our paper. In [MN12], D. Niwinski and H. Michalewski are also interested in the problem of finding upper bounds for the class of regular tree languages. Using a method developed by J. Saint Raymond in [SR06], they prove that the game tree language $W_{1,3}$ is complete for the class of Σ_1^1 -inductive sets. The Σ_1^1 -inductive sets are known to contain more complex sets than the σ -algebra generated by the analytic sets. Moreover, the referee indicated us that one can actually verify that the language given in S. Hummel’s paper [Hum12] is reducible to $W_{1,3}$ and is not complete for the class of Σ_1^1 -inductive sets. In another recent paper [GMMS14] T. Gogacz, H. Michalewski, M. Mio and M. Skrzypczak show a one-to-one correspondence between the levels of the hierarchy of Kolmogorov R-sets and parity index of regular languages which extends the theorem from [MN12]. Since due to a theorem of Burgess the R-sets are known to be in correspondence with the game quantifier, on a technical level this covers Theorem 4.4. We also notice that another estimation of the kind of Theorem 5.13 is present in [GMMS14], where it is shown that regular languages of infinite trees occupy exactly the first ω -levels of Kolmogorov’s hierarchy; and on the other hand it is known that there exist ω_1 levels in this hierarchy, all of them contained in the class Δ_2^1 . Theorem 5.13 may then be seen as a generalization of the estimation following from [GMMS14].

We now state some important open questions.

- A difficult problem in the study of the topological complexity of recognizable tree languages is to determine the Wadge hierarchy of tree languages accepted by non deterministic Muller or Rabin tree automata. A subquestion is to know whether there is a regular language of infinite trees which is a Borel set of *infinite rank*.
- A related problem is to study the determinacy of Wadge games between tree automata; this would be a first step towards the possibility of using these games to determine the topological complexity of a regular tree language. Note that in the case of one-counter Büchi automata it has been shown that the determinacy of such games needs some large cardinal assumption (see [Fin13a]).

We now state some important open questions which are more related to the computer science roots of regular languages since they also involve decidability questions.

- While many questions about *deterministic* regular languages of infinite trees have been shown to be decidable [ADMN07], many corresponding questions about *non-deterministic* regular languages of infinite trees are still open.

Concerning the decidability of the topological complexity of regular languages, it has been shown recently by M. Bojanczyk and T. Place in [BP12] that one can decide whether a regular language accepted by a given tree automaton is a boolean combination of open sets. This result has been extended by A. Facchini and H. Michalewski in [FM14], where the authors prove that one can decide whether a regular tree language is in the class Δ_2^0 . The question is still open for the other levels of the Borel hierarchy, or whether a regular tree language is Borel, analytic, coanalytic, or in any class $\Sigma\Gamma$ present in this paper.

Acknowledgement. We wish to thank the anonymous referee for very useful comments on a preliminary version of this paper which lead to a great improvement of the paper, and for indicating us some related studies that are now present in the bibliography.

References

- [ADMN07] A. Arnold, J. Duparc, F. Murlak, and D. Niwinski. On the topological complexity of tree languages. In J. Flum, E. Grädel, and T. Wilke, editors, *Logic and Automata: History and Perspectives*, pages 9–28. Amsterdam University Press, 2007.
- [AN08] A. Arnold and D. Niwinski. Continuous separation of game languages. *Fundamenta Informaticae*, 81(1–3):19–28, 2008.
- [BDQ05] J. C. Bradfield, J. Duparc, and S. Quickert. Transfinite extension of the mu-calculus. In C.-H. Luke Ong, editor, *Computer Science Logic, 19th International Workshop, CSL 2005, 14th Annual Conference of the EACSL, Oxford, UK, August 22-25, 2005, Proceedings*, volume 3634 of *Lecture Notes in Computer Science*, pages 384–396. Springer, 2005.
- [BP12] M. Bojanczyk and T. Place. Regular languages of infinite trees that are boolean combinations of open sets. In Artur Czumaj, Kurt Mehlhorn, Andrew M. Pitts, and Roger Wattenhofer, editors, *Automata, Languages, and Programming - 39th International Colloquium, ICALP 2012, Warwick, UK, July 9-13, 2012, Proceedings, Part II*, volume 7392 of *Lecture Notes in Computer Science*, pages 104–115. Springer, 2012.
- [Bra03] J. C. Bradfield. Fixpoints, games and the difference hierarchy. *RAIRO-Theoretical Informatics and Applications*, 37(1):1–15, 2003.
- [CS07] B. Cagnard and P. Simonnet. Baire and automata. *Discrete Mathematics and Theoretical Computer Science*, 9(2):255–296, 2007.

- [Dup01] J. Duparc. Wadge hierarchy and Veblen hierarchy: Part 1: Borel sets of finite rank. *Journal of Symbolic Logic*, 66(1):56–86, 2001.
- [Fin13a] O. Finkel. The determinacy of context-free games. *The Journal of Symbolic Logic*, 78(4):1115–1134, 2013.
- [Fin13b] O. Finkel. Infinite games specified by 2-tape automata. 2013. Preprint, available from <http://fr.arxiv.org/abs/1312.3797>.
- [FM14] A. Facchini and H. Michalewski. Deciding the Borel complexity of regular tree languages. In Arnold Beckmann, Erzsébet Csuhaj-Varjú, and Klaus Meer, editors, *Language, Life, Limits - 10th Conference on Computability in Europe, CiE 2014, Budapest, Hungary, June 23-27, 2014. Proceedings*, volume 8493 of *Lecture Notes in Computer Science*, pages 163–172. Springer, 2014.
- [FS09] O. Finkel and P. Simonnet. On recognizable tree languages beyond the Borel hierarchy. *Fundamenta Informaticae*, 95(2-3):287–303, 2009.
- [GH82] Y. Gurevich and L. Harrington. Trees, automata, and games. In Harry R. Lewis, Barbara B. Simons, Walter A. Burkhard, and Lawrence H. Landweber, editors, *Proceedings of the 14th Annual ACM Symposium on Theory of Computing, May 5-7, 1982, San Francisco, California, USA*, pages 60–65. ACM, 1982.
- [GMMS14] T. Gogacz, H. Michalewski, M. Mio, and M. Skrzypczak. Measure properties of game tree languages. In Erzsébet Csuhaj-Varjú, Martin Dietzfelbinger, and Zoltán Ésik, editors, *Mathematical Foundations of Computer Science 2014 - 39th International Symposium, MFCS 2014, Budapest, Hungary, August 25-29, 2014. Proceedings, Part I*, volume 8634 of *Lecture Notes in Computer Science*, pages 303–314. Springer, 2014.
- [GTW02] E. Grädel, W. Thomas, and W. Wilke, editors. *Automata, Logics, and Infinite Games: A Guide to Current Research [outcome of a Dagstuhl seminar, February 2001]*, volume 2500 of *Lecture Notes in Computer Science*. Springer, 2002.
- [HKMN08] G. Hjorth, B. Khoussainov, A. Montalbán, and A. Nies. From automatic structures to Borel structures. In *Proceedings of the Twenty-Third Annual IEEE Symposium on Logic in Computer Science, LICS 2008, 24-27 June 2008, Pittsburgh, PA, USA*, pages 431–441. IEEE Computer Society, 2008.

- [Hum12] S. Hummel. Unambiguous tree languages are topologically harder than deterministic ones. In Marco Faella and Aniello Murano, editors, *Proceedings Third International Symposium on Games, Automata Logics and Formal Verification, GandALF 2012, Napoli, Italy, September 6-8, 2012*, volume 96 of *EPTCS*, pages 247–260, 2012.
- [Jec02] T. Jech. *Set theory, third edition*. Springer, 2002.
- [Kan97] A. Kanamori. *The Higher Infinite*. Springer-Verlag, 1997.
- [Kec95] A. S. Kechris. *Classical descriptive set theory*. Springer-Verlag, New York, 1995.
- [LSR88] A. Louveau and J. Saint-Raymond. The strength of Borel Wadge determinacy. In *Cabal Seminar 81–85*, volume 1333 of *Lecture Notes in Mathematics*, pages 1–30. Springer, 1988.
- [LT94] H. Lescow and W. Thomas. Logical specifications of infinite computations. In J. W. de Bakker, Willem P. de Roever, and Grzegorz Rozenberg, editors, *A Decade of Concurrency*, volume 803 of *Lecture Notes in Computer Science*, pages 583–621. Springer, 1994.
- [MN12] H. Michalewski and D. Niwinski. On topological completeness of regular tree languages. In Robert L. Constable and Alexandra Silva, editors, *Logic and Program Semantics - Essays Dedicated to Dexter Kozen on the Occasion of His 60th Birthday*, volume 7230 of *Lecture Notes in Computer Science*, pages 165–179. Springer, 2012.
- [Mos09] Y. N. Moschovakis. *Descriptive set theory*, volume 155 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, second edition, 2009.
- [Mur08] F. Murlak. The Wadge hierarchy of deterministic tree languages. *Logical Methods in Computer Science*, 4(4, paper 15), 2008.
- [Niw85] D. Niwinski. An example of non Borel set of infinite trees recognizable by a Rabin automaton. 1985. in Polish, manuscript.
- [NW03] D. Niwinski and I. Walukiewicz. A gap property of deterministic tree languages. *Theoretical Computer Science*, 1(303):215–231, 2003.
- [PP04] D. Perrin and J.-E. Pin. *Infinite words, automata, semigroups, logic and games*, volume 141 of *Pure and Applied Mathematics*. Elsevier, 2004.

- [Rab69] M. O. Rabin. Decidability of second-order theories and automata on infinite trees. *Transactions of the American Mathematical Society*, 141:1–35, 1969.
- [Sim92] P. Simonnet. *Automates et théorie descriptive*. PhD thesis, Université Paris VII, 1992.
- [Sku93] J. Skurczynski. The Borel hierarchy is infinite in the class of regular sets of trees. *Theoretical Computer Science*, 112(2):413–418, 1993.
- [SR06] J. Saint Raymond. Quasi-bounded trees and analytic inductions. *Fundamenta Mathematicae*, 191(2):175–185, 2006.
- [Sta97] L. Staiger. ω -languages. In *Handbook of formal languages, Vol. 3*, pages 339–387. Springer, Berlin, 1997.
- [Tho90] W. Thomas. Automata on infinite objects. In J. van Leeuwen, editor, *Handbook of Theoretical Computer Science*, volume B, Formal models and semantics, pages 135–191. Elsevier, 1990.
- [Tho97] W. Thomas. Languages, automata, and logic. In *Handbook of formal languages, Vol. 3*, pages 389–455. Springer, Berlin, 1997.
- [Wad83] W. Wadge. *Reducibility and determinateness in the Baire space*. PhD thesis, University of California, Berkeley, 1983.